LAMPRECHT-TATE FORMULA

SAZZAD ALI BISWAS

Abstract. For multiplicative character of a non-Archimedean local field, we have a formula for epsilon factor due to John Tate. Before Tate, Erich Lamprecht, also gave a formula for local epsilon factors of linear characters. In this paper, we generalize the Tate's formula and show that the Lamprecht and Tate formulas are the special cases of our formula. We also show that the famous twisting formula of Deligne is a corollary of our formula.

arXiv:1702.04286v1 [math.NT] 14 Feb 2017

1. Introduction

By Langlands we can associate a local epsilon factor (also known as local constant) with each finite dimensional continous complex representation ρ of the absolute Galois group G_F of a non-Archimedean local field F. But there is no explicit formula for local epsilon of any arbitrary local Galois representation. In 1952 [6], Erich Lamprecht gave a formula (cf. Lemma 8.1 on p. 60 of [7]) for local epsilon factor of a linear character $\chi : F^\times \to \mathbb{C}^\times$. Later, John Tate, gave more explicit formula (cf. [9], p. 94) for local epsilon factor of linear character. In this paper, we generalize the Tate's formula (2.6) and give Theorem 3.1. We also show that the Tate's formula (2.6) and Lamprecht formula (cf. Corollary 3.2(1)) are the special cases of our formula (3.2).

Furthermore, we also show that the famous Deligne's twisting formula (cf. Corollary 3.2(2)) of local epsilon factors is a corollary of the Theorem 3.1.

2. Notations and Preliminaries

Let F be a non-Archimedean local field of characteristic zero , i.e., a finite extension of the field $\mathbb{Q}_p$ (field of p-adic numbers), where p is a prime. Let O_F be the ring of integers in local field F and $P_F = \pi_F O_F$ is the unique prime ideal in O_F and π_F is a uniformizer, i.e., an element in P_F whose valuation is one, i.e., $\nu_F(\pi_F) = 1$. The cardinality of the residue field $\mathbb{F}_q = O_F/P_F$ of F is q_F, i.e., $|\mathbb{F}_q| = q_F$. Let $U_F = O_F - P_F$ be the group of units in O_F. Let $P_F^i = \{x \in F : \nu_F(x) \geqslant i\}$ and for $i \geqslant 0$ define $U_F^i = 1 + P_F^i$ (with proviso $U_F^0 = U_F = O_F^\times$).

Definition 2.1 (**Different and Discriminant**)**.** Let K/F be a finite separable extension of non-Archimedean local field F. We define the **inverse different (or codifferent)** $\mathcal{D}_{K/F}^{-1}$ of K over F to be $\pi_K^{-d_{K/F}} O_K$, where $d_{K/F}$ is the largest integer (this is the exponent of the different $\mathcal{D}_{K/F}$) such that

$$\mathrm{Tr}_{K/F}(\pi_K^{-d_{K/F}} O_K) \subseteq O_F,$$

where $\mathrm{Tr_{K/F}}$ is the trace map from K to F. Then the **different** is defined by:

$$\mathcal{D}_{K/F} = \pi_K^{d_{K/F}} O_K$$

and the **discriminant** $D_{K/F}$ is

2010 *Mathematics Subject Classification.* 11S37 (11F70, 22E50).

Key words and phrases. Local fields, Characters, Epsilon factors, Conductor.

The author is partially supported by Infosys Foundation, India .

$$D_{K/F} = N_{K/F}(\pi_K^{d_{K/F}})O_F.$$

Definition 2.2 (**Conductor of characters**)**.** The conductor of any nontrivial additive character ψ of a field is an integer $n(\psi)$ if ψ is trivial on $P_F^{-n(\psi)}$, but nontrivial on $P_F^{-n(\psi)-1}$. We also consider $a(\chi)$ as the conductor of nontrivial character $\chi : F^\times \to \mathbb{C}^\times$, i.e., $a(\chi)$ is the smallest integer $m \geqslant 0$ such that χ is trivial on U_F^m. We say χ is **unramified** if the conductor of χ is **zero** and otherwise **ramified**. We also recall here that for two characters χ_1 and χ_2 of $F^\times$ we have $a(\chi_1\chi_2) \leqslant \max(a(\chi_1), a(\chi_2))$ with equality if $a(\chi_1) \neq a(\chi_2)$.

2.1. **Classical Gauss sums.** Let k_q be a finite field. Let p be the characteristic of k_q; then the prime field contained in k_q is k_p. The structure of the **canonical** additive character ψ_q of k_q is the same as the structure of the canonical character ψ_F, namely **it comes by trace** from the canonical character of the base field, i.e.,

$$\psi_q = \psi_p \circ \mathrm{Tr}_{\mathrm{k_q/k_p}},$$

where

$$\psi_p(x) := e^{\frac{2\pi i x}{p}} \qquad \text{for all } x \in k_p.$$

Gauss sums: Let χ be a multiplicative and ψ an additive character of k_q. Then the Gauss sum $G(\chi, \psi)$ is define by

$$G(\chi, \psi) = \sum_{x \in k_q^\times} \chi(x)\psi(x). \tag{2.1}$$

In the following theorem for quadratic characters of k_q we can give explicit formula of Gauss sums.

Theorem 2.3 ([8], p. 199, Theorem 5.15)**.** *Let k_q be a finite field with $q = p^s$, where p is an odd prime and $s \in \mathbb{N}$. Let χ be the quadratic character of k_q and let ψ be the canonical additive character of k_q. Then*

$$G(\chi, \psi) = \begin{cases} (-1)^{s-1} q^{\frac{1}{2}} & \text{if } p \equiv 1 \pmod 4, \\ (-1)^{s-1} i^s q^{\frac{1}{2}} & \text{if } p \equiv 3 \pmod 4. \end{cases} \tag{2.2}$$

2.2. **Epsilon factors.** For a nontrivial multiplicative character χ of $F^\times$ and nontrivial additive character ψ of F, we have (cf. [7], p. 4)

$$\epsilon(\chi, \psi, c) = \chi(c) \frac{\int_{U_F} \chi^{-1}(x)\psi(x/c)dx}{|\int_{U_F} \chi^{-1}(x)\psi(x/c)dx|} \tag{2.3}$$

where the Haar measure dx on F is normalized such that the Haar measure m' of O_F is 1, i.e., $m'(O_F) = 1$, and $c \in F^\times$ with valuation $n(\psi) + a(\chi)$. To get modified **summation formula** of epsilon factor from the above integral formula (2.3), we need the next lemma which can be found in [4].

Lemma 2.4. *Let F be a non-Archimedean local field with q_F as the cardinality of the residue field of F. Let χ be a nontrivial character of $F^\times$ with conductor $a(\chi)$. Let ψ be an additive character of F with conductor $n(\psi)$. We define the integration for an integer $m \in \mathbb{Z}$:*

$$I(m) = \int_{U_F} \chi^{-1}(x)\psi(\frac{x}{\pi_F^{l+m}})dx, \quad \text{where } l = a(\chi) + n(\psi). \tag{2.4}$$

Then

$$|I(m)| = \begin{cases} q_F^{-\frac{a(\chi)}{2}} & \text{when } m = 0 \\ 0 & \text{otherwise} \end{cases} \tag{2.5}$$

Again since $m'(O_F) = 1$, we have $m'(U_F^n) = q_F^{-n}$ for any positive integer n. By using the above Lemma 2.4, the formula (2.3) can be reduced to (cf. p. 94 of [9])

$$\begin{aligned} \epsilon(\chi, \psi) &= \chi(c) q_F^{\frac{a(\chi)}{2}} \sum_{x \in \frac{U_F}{U_F^{a(\chi)}}} \chi^{-1}(x)\psi(x/c) m'(U_F^{a(\chi)}) \\ &= \chi(c) q_F^{-\frac{a(\chi)}{2}} \sum_{x \in \frac{U_F}{U_F^{a(\chi)}}} \chi^{-1}(x)\psi(x/c), \end{aligned} \tag{2.6}$$

where $c = \pi_F^{a(\chi)+n(\psi)}$.

3. **Lamprecht-Tate formula for epsilon factor**

Theorem 3.1 (**Lamprecht-Tate formula**). *Let F be a non-Archimedean local field. Let χ be a character of $F^\times$ of conductor $a(\chi)$ and let m be a natural number such that $2m \leqslant a(\chi)$. Let ψ_F be the canonical additive character of F. Then there exists $c \in F^\times$, $\nu_F(c) = a(\chi) + d_{F/\mathbb{Q}_p}$ such that*

$$\chi(1+y) = \psi_F(c^{-1}y) \qquad \text{for all } y \in P_F^{a(\chi)-m}, \tag{3.1}$$

and this will imply:

$$\epsilon(\chi, \psi_F) = \chi(c) \cdot q_F^{-\frac{(a(\chi)-2m)}{2}} \sum_{x \in (1+P_F^m)/(1+P_F^{a(\chi)-m})} \chi^{-1}(x)\psi_F(c^{-1}x). \tag{3.2}$$

Remark: *Note that the assumption (3.1) is obviously fulfilled for $m = 0$ because then both sides are $= 1$, and the resulting formula for $m = 0$ is the Tate formula (2.6).*

Proof. When $m = 0$, the formula (3.2) is the same as the Tate's formula (2.6), hence for $m = 0$ everything is correct. In general, the assumption $2m \leqslant a(\chi)$ implies $2(a(\chi) - m) \geqslant a(\chi)$ and therefore

$$\chi(1+y)\chi(1+y') = \chi(1+y+y')$$

for $y, y' \in P_F^{a(\chi)-m}$. That is, $y \mapsto \chi(1+y)$ is a character of the additive group $P_F^{a(\chi)-m}$. This character extends to a character of F^+ and, by local additive duality, there is some $c \in F^\times$ such that

$$\chi(1+y) = \psi_F(c^{-1}y) = (c^{-1}\psi_F)(y), \quad \text{for all } y \in P_F^{a(\chi)-m}.$$

Now comparing the conductors of both sides we must have:

$$a(\chi) = -n(c^{-1}\psi_F) = \nu_F(c) - n(\psi_F),$$

hence $\nu_F(c) = a(\chi) + n(\psi_F) = a(\chi) + d_{F/\mathbb{Q}_p}$ is the right assumption for our formula.
Now we assume $m \geqslant 1$ (the case $m = 0$ we have checked already) and consider the filtration

$$O_F^\times \supseteq 1 + P_F^{a(\chi)-m} \supseteq 1 + P_F^{a(\chi)}.$$

Then we may represent $x \in O_F^\times/(1+P_F^{a(\chi)})$ as $x = z(1+y)$, where $y \in P_F^{a(\chi)-m}$ and z runs over the system of representatives for $O_F^\times/(1+P_F^{a(\chi)-m})$. Now computing $\epsilon(\chi,\psi_F)$ we have to consider the sum

(3.3)
$$\sum_{x\in O_F^\times/(1+P_F^{a(\chi)})} \chi^{-1}(x)\psi_F(c^{-1}x) = \sum_{z\in O_F^\times/(1+P_F^{a(\chi)-m})} \sum_{y\in P_F^{a(\chi)-m}/P_F^{a(\chi)}} \chi^{-1}(z(1+y))\psi_F(c^{-1}z(1+y)).$$

Now using (3.1) we obtain

$$\chi^{-1}(z(1+y)) = \chi^{-1}(z)\chi^{-1}(1+y) = \chi^{-1}(z)\chi(1-y) = \chi^{-1}(z)\psi_F(-c^{-1}y)$$

and therefore our double sum (3.3) rewrites as

$$\sum_{z\in O_F^\times/(1+P_F^{a(\chi)-m})} \chi^{-1}(z)\psi_F(c^{-1}z)\cdot\left(\sum_{y\in P_F^{a(\chi)-m}/P_F^{a(\chi)}} \psi_F(c^{-1}y(z-1))\right).$$

But the inner sum is the sum on the additive group $P_F^{a(\chi)-m}/P_F^{a(\chi)}$ and $(c^{-1}(z-1))\psi_F$ is a character of that group. Hence this sum is equal to $[P_F^{a(\chi)-m} : P_F^{a(\chi)}] = q_F^m$ if the character is $\equiv 1$ and otherwise the sum will be zero. But:

$$n(c^{-1}(z-1)\psi_F) = \nu_F(c^{-1}(z-1)) + n(\psi_F) = -a(\chi) + \nu_F(z-1).$$

So the character $(c^{-1}(z-1))\psi_F$ is trivial on $P_F^{a(\chi)-\nu_F(z-1)}$, and therefore it will be $\equiv 1$ on $P_F^{a(\chi)-m}$ if and only if $\nu_F(z-1) \geqslant m$, i.e., $z = 1+y' \in 1+P_F^m$. Therefore our sum (3.3) rewrites as

(3.4)
$$\sum_{x\in O_F^\times/(1+P_F^{a(\chi)})} \chi^{-1}(x)\psi_F(c^{-1}x) = q_F^m \sum_{z\in(1+P_F^m)/(1+P_F^{a(\chi)-m})} \chi^{-1}(z)\psi_F(c^{-1}z).$$

And substituting this result into the Tate formula (2.6) we get

$$\begin{aligned}\epsilon(\chi,\psi_F) &= \chi(c)q_F^{-\frac{a(\chi)}{2}} \sum_{x\in O_F^\times/(1+P_F^{a(\chi)})} \chi^{-1}(x)\psi_F(c^{-1}x)\\ &= \chi(c)\cdot q_F^{-\frac{(a(\chi)-2m)}{2}} \sum_{x\in(1+P_F^m)/(1+P_F^{a(\chi)-m})} \chi^{-1}(x)\psi_F(c^{-1}x).\end{aligned}$$

□

Corollary 3.2. *(1)* **Lamprecht Formulas (cf. Lemma 8.1, p.60 of** [7]**)** *Let χ be a character of $F^\times$. Let ψ be a nontrivial additive character of F.*

(a) When $a(\chi) = 2d\,(d \geqslant 1)$, we have

$$\epsilon(\chi,\psi) = \chi(c)\psi(c^{-1}).$$

(b) When $a(\chi) = 2d+1\,(d \geqslant 1)$, we have

$$\epsilon(\chi,\psi) = \chi(c)\psi(c^{-1})\cdot q_F^{-\frac{1}{2}} \sum_{x\in P_F^d/P_F^{d+1}} \chi^{-1}(1+x)\psi(c^{-1}x).$$

(2) **Deligne's twisting formula (cf.** [3]**, Lemma 4.16):** *If $\alpha,\beta \in \widehat{F^\times}$ with $a(\alpha) \geqslant 2\cdot a(\beta)$, then*

(3.5)
$$\epsilon(\alpha\beta,\psi) = \beta(c)\cdot\epsilon(\alpha,\psi).$$

Here $c \in F^\times$ with F-valuation $\nu_F(c) = a(\chi) + n(\psi_F)$, and in Case (1(a)) and Case (1(b)), c also satisfies

$$\chi(1+x) = \psi(\tfrac{x}{c}) \text{ for all } x \in F^\times \text{ with } 2 \cdot \nu_F(x) \geqslant a(\chi).$$

And in case (2), we have $\alpha(1+x) = \psi(x/c)$ for all $\nu_F(x) \geqslant \frac{a(\alpha)}{2}$

Proof. From the above formula (3.2), the assertions are followed.
(1(a)). When $a(\chi) = 2d$, where $d \geqslant 1$. In this case, we take $m = d$, and from equation (3.2) we obtain

$$\epsilon(\chi, \psi) = \chi(c) \cdot \sum_{x \in (1+P_F^d)/(1+P_F^d)} \chi^{-1}(x)\psi(c^{-1}x) = \chi(c) \cdot \psi(c^{-1}). \tag{3.6}$$

(1(b)). When $a(\chi) = 2d+1$, where $d \geqslant 1$. In this case, we also take $m = d$, and then from equation (3.2) we obtain

$$\begin{aligned}\epsilon(\chi, \psi) &= \chi(c) \cdot q_F^{-\frac{1}{2}} \cdot \sum_{x \in (1+P_F^d)/(1+P_F^{d+1})} \chi^{-1}(x)\psi(c^{-1}x) \\ &= \chi(c) \cdot \psi(c^{-1}) \cdot q_F^{-\frac{1}{2}} \cdot \sum_{x \in P_F^d/P_F^{d+1}} \chi^{-1}(1+x)\psi(c^{-1}x).\end{aligned}$$

Here $c \in F^\times$ with $\nu_F(c) = a(\chi) + n(\psi)$, satisfies

$$\chi(1+x) = \psi(c^{-1}x) \text{ for all } x \in F^\times \text{ with } 2\nu_F(x) \geqslant a(\chi).$$

(2) Proof of Deligne's twisting formula: By the given condition, $a(\alpha) \geqslant 2a(\beta)$, hence we have $a(\alpha\beta) = a(\alpha)$. Now take $m = a(\beta)$, then from equation (3.2) we can write:

$$\begin{aligned}\epsilon(\alpha\beta, \psi) &= \alpha\beta(c) \cdot q_F^{-\frac{(a(\alpha)-2m)}{2}} \cdot \sum_{x \in (1+P_F^m)/(1+P_F^{a(\chi)-m})} (\alpha\beta)^{-1}(x)\psi(\frac{x}{c}) \\ &= \beta(c) \cdot \alpha(c) q_F^{-\frac{(a(\alpha)-2m)}{2}} \sum_{x \in (1+P_F^m)/(1+P_F^{a(\alpha)-m})} \alpha^{-1}(x)\psi(\frac{x}{c}) \\ &= \beta(c) \cdot \epsilon(\alpha, \psi),\end{aligned}$$

since $a(\beta) = m$, hence $\beta(x) = 1$ for all $x \in (1+P_F^m)/(1+P_F^{a(\alpha)-m})$.

□

Remark 3.3. Let μ_{p^∞} denote as the group of roots of unity of p-power order.

Now we can give an explicit formula for $\epsilon(\chi, \psi_F)$ modulo μ_{p^∞} (cf. the review of Henniart's paper [5] by E.-W. Zink) and which is:

(1) $\epsilon(\chi, \psi_F) \equiv \chi(c) \mod \mu_{p^\infty}$ if $a(\chi)$ is even,
(2) $\epsilon(\chi, \psi_F) \equiv \chi(c)G(c) \mod \mu_{p^\infty}$ if $a(\chi) = 2d+1$, where

$$G(c) := q_F^{-\frac{1}{2}} \cdot \sum_{x \in P_F^d/P_F^{d+1}} \psi_F(\frac{x^2}{2c})$$

depends only on $c \in F^\times \mod 1 + P_F$.

Remark 3.4 (**On** $G(c)$)**.** Let $\mathbb{F}_q$ be a finite field with q element and q is odd. Let ψ_0 be a nontrivial additive character of $\mathbb{F}_q$. Let η be the quadratic character of $\mathbb{F}_q^\times$. Then from the generalization (cf. [1], p. 47, Problem 24) of Theorem 1.1.5 of [1] on p. 12 we have

$$\sum_{x\in\mathbb{F}_q}\psi_0(x^2)=\sum_{x\in\mathbb{F}_q^\times}\eta(x)\psi_0(x)=G(\eta,\psi_0). \tag{3.7}$$

Now from Theorem 2.3 we can observe that $q^{-\frac{1}{2}}\sum_{x\in\mathbb{F}_q}\psi_0(x^2)$ is a **fourth** root of unity.

Since $\nu_F(c)=a(\chi)+n(\psi_F)$, $P_F^d/P_F^{d+1}\cong k_F$, we can deduce $G(c)$ as

$$G(c)=q_F^{-\frac{1}{2}}\sum_{x\in k_F}\psi'(x^2)=q_F^{-\frac{1}{2}}G(\eta,\psi'),$$

where ψ' is a nontrivial additive character of k_F and η is the quadratic character of $k_F^\times$. Therefore we can conclude that $G(c)$ is a fourth root of unity.

Furthermore, since $G(c)\cdot q_F^{\frac{1}{2}}=G(\eta,\psi')$ is a quadratic classical Gauss sum, from the computation of lambda function $\lambda_{K/F}(\psi)$ (for definition of lambda functions, see equation (3.2.7) on p. 37 of [2]), where K/F is a tamely ramified quadratic extension (cf. Subsection 3.4.1 of [2]), we can conclude that $G(c)$ is actually a λ-function for some quadratic tamely ramified local extension.

Acknowledgements I would like to thank Professor E.-W. Zink for encouraging me to work on local epsilon factors and his constant valuable advices.

References

[1] B.C. Berndt, R.J. Evans, K.S. Williams, Gauss and Jacobi Sums, John Wiley and Sons, INC,

[2] S.A. Biswas, Local constants for Galois representations - Some explicit results, Ph.D Thesis, `https://arxiv.org/pdf/1603.06089.pdf`.

[3] P. Deligne, Les constantes des équations fonctionnelle des fonctions L, in Modular functions of one variable II, Lecture Notes in Mathematics **349**(1972), 501-597, Springer-Verlag, Berlin-Heidelberg-New York.

[4] I.M. Gelfand, M.I. Graev, I.I. Pyatetski-Shapiro, Representation theory and automorphic functions, W.B. Saunders Company, 1969.

[5] G. Henniart, Galois ϵ-factors modulo roots of unity, Invent. Math. **78** (1984), 117-126.

[6] E. Lamprecht, Allgemeine theorie der Gauss'schen summen in endlichen kommutativen ringen, Math. Nach. **9**(1953).

[7] R. P. Langlands, On the functional equation of the Artin L-functions, unpublished article, `https://publications.ias.edu/sites/default/files/a-ps.pdf`.

[8] R. Lidl, H. Niederreiter, Finite fields, Encyclopedia of Mathematics and its applications, Cambridge University press 2000.

[9] J. Tate, Local Constants, Algebraic Number Fields (L-functions and Galois properties), Proceedings of Symposium, Edited by A. Fröhlich, pp. 89-131.

Chennai Mathematical Institute, H1, Sipcot It Park, Siruseri, Kelambakkam, 603103, India
E-mail address: `sabiswas@cmi.ac.in, sazzad.jumath@gmail.com`